

## List of Sessions of Interest:

Vepdegestrant, a PROTAC estrogen receptor (ER) degrader, vs fulvestrant in ER-positive/human epidermal growth factor receptor 2 (HER2)-negative advanced breast cancer: Results of the global, randomized, phase 3 VERITAC-2 study.
Phase 3 trial of adjuvant cemiplimab (cemi) versus placebo (pbo) for high-risk cutaneous squamous cell carcinoma (CSCC).
Trastuzumab deruxtecan (T-DXd) vs ramucirumab (RAM) + paclitaxel (PTX) in second-line treatment of patients (pts) with human epidermal growth factor receptor 2-positive (HER2+) unresectable/metastatic gastric cancer (GC) or gastroesophageal junction adenocarcinoma (GEJA): Primary analysis of the randomized, phase 3 DESTINY-Gastric04 study.
TROPION-Lung02: Datopotamab deruxtecan (Dato-DXd) plus pembrolizumab (pembro) with or without platinum chemotherapy (Pt-CT) as first-line (1L) therapy for advanced non-small cell lung cancer (aNSCLC).
Randomized trial of standard chemotherapy alone or combined with atezolizumab as adjuvant therapy for patients with stage III deficient DNA mismatch repair (dMMR) colon cancer (Alliance A021502; ATOMIC).
NIVOPOSTOP (GORTEC 2018-01): A phase III randomized trial of adjuvant nivolumab added to radio-chemotherapy in patients with resected head and neck squamous cell carcinoma at high risk of relapse.
Camizestran + CDK4/6 inhibitor (CDK4/6i) for the treatment of emergent ESR1 mutations during first-line (1L) endocrine-based therapy (ET) and ahead of disease progression in patients (pts) with HR+/HER2- advanced breast cancer (ABC): Phase 3, double-blind ctDNA-guided SERENA-6 trial.
Event-free survival (EFS) in MATTERHORN: A randomized, phase 3 study of durvalumab plus 5-fluorouracil, leucovorin, oxaliplatin and docetaxel chemotherapy (FLOT) in resectable gastric/gastroesophageal junction cancer (GC/GEJC).
FIRST/ENGOT-OV44: A phase 3 clinical trial of dostarlimab (dost) and niraparib (nira) in first-line (1L) advanced ovarian cancer (aOC).
ROSELLA: A phase 3 study of relacorilant in combination with nab-paclitaxel versus nab-paclitaxel monotherapy in patients with platinum-resistant ovarian cancer (GOG-3073, ENGOT-ov72).
Linvoseltamab (LINV0) + carfilzomib (CFZ) in patients (pts) with relapsed/refractory multiple myeloma (RRMM): Initial results from the LINKER-MM2 trial.
Neoadjuvant (neoadj) osimertinib (osi) ± chemotherapy (CT) vs CT alone in resectable (R) epidermal growth factor receptor-mutated (EGFRm) NSCLC: NeoADAURA.
Lurbinectedin (lurbi) + atezolizumab (atezo) as first-line (1L) maintenance treatment (tx) in patients (pts) with extensive-stage small cell lung cancer (ES-SCLC): Primary results of the phase 3 IMforte trial.
Tarlatamab versus chemotherapy (CTx) as second-line (2L) treatment for small cell lung cancer (SCLC): Primary analysis of Ph3 DeLLphi-304.
Glofitamab plus gemcitabine and oxaliplatin (Glofit-GemOx) in patients (pts) with relapsed/refractory (R/R) diffuse large B-cell lymphoma (DLBCL): 2-year (yr) follow-up of STARGLO.
INAVO120: Phase III trial final overall survival (OS) analysis of first-line inavolisib (INAVO)/placebo (PBO) + palbociclib (PALBO) + fulvestrant (FULV) in patients (pts) with PIK3CA-mutated, hormone receptor-positive (HR+), HER2-negative (HER2-), endocrine-resistant advanced breast cancer (aBC).
Sasanlimab in combination with bacillus Calmette-Guérin (BCG) in BCG-naïve, high-risk non-muscle-invasive bladder cancer (NMIBC): Event-free survival (EFS) subgroup analyses based on disease stage from the CREST study.
Mevrometostat in combination with enzalutamide in patients with metastatic castration-resistant prostate cancer (mCRPC) previously treated with abiraterone acetate: The phase 3, randomized MEVPRO-1 study.
Results from VERIFY, a phase 3, double-blind, placebo (PBO)-controlled study of rusfertide for treatment of polycythemia vera (PV).

Sigvotatug vedotin (SV), an investigational integrin beta-6 (IB6)-directed antibody–drug conjugate (ADC), and pembrolizumab combination therapy: Initial results from an ongoing phase 1 study (SGNB6A-001).
First-line encorafenib + cetuximab + mFOLFOX6 in BRAF V600E-mutant metastatic colorectal cancer (BREAKWATER): Progression-free survival and updated overall survival analyses.
SOHO-01: Safety and efficacy of BAY 2927088 in patients with advanced HER2-mutant non-small cell lung cancer (NSCLC) who were pretreated but naïve to HER2-targeted therapy or had not received any treatment for advanced disease.
Sasanlimab as bladder-sparing maintenance treatment after neoadjuvant chemotherapy in patients with muscle invasive bladder cancer (MIBC): The phase 2, SASAN-SPARING trial.
Sacituzumab govitecan (SG) + pembrolizumab (pembro) vs chemotherapy (chemo) + pembro in previously untreated PD-L1–positive advanced triple-negative breast cancer (TNBC): Primary results from the randomized phase 3 ASCENT-04/KEYNOTE-D19 study.
Trastuzumab deruxtecan (T-DXd) + pertuzumab (P) vs taxane + trastuzumab + pertuzumab (THP) for first-line (1L) treatment of patients (pts) with human epidermal growth factor receptor 2–positive (HER2+) advanced/metastatic breast cancer (a/mBC): Interim results from DESTINY-Breast09.
Safety and efficacy of chimeric antigen receptor (CAR)–T cell therapy (BRG01) targeting the Epstein-Barr virus (EBV) envelope protein in EBV+ lymphoproliferative disease patients.